

Mixed Features in Bipolar I Disorder


Introduction

Bipolar I disorder (BP-1) is a mood disorder characterized by alternating states of opposing mood polarity, depressive and manic or hypomanic episodes. While the transition between the mood states is a hallmark feature of bipolar disorder (BP),¹ this bipolarity of mood can exist in a spectrum with varying degrees of severity.² Mood symptoms of opposing polarity, depressive and manic features, can occur simultaneously at subthreshold levels in what is called a “mixed state.”³

Prevalence and Impact of Mixed Features in BP Patients

Mixed moods are increasingly being recognized as a significant component of BP,⁴ as a number of bipolar patients may experience mixed episodes. A study showed that approximately 30% to 34% of patients with BP experience mixed features as defined by the *Diagnostic and Statistical Manual of Mental Disorders (DSM-5)*.³ In another study, 34% of patients with BP-1 and 26% of patients with major depressive disorder (MDD) were shown to exhibit mixed features during a depressive episode. Additionally, 20% of BP-1 patients exhibited mixed features during a manic/hypomanic episode.⁵

30%-34%
of patients with
bipolar disorder experience
mixed features



Mixed features in BP-1 may be associated with more frequent episodes and a more severe course of illness. Studies have found that bipolar patients with mixed features can experience^{5,9-11}:

- A longer duration of episodes
- Higher likelihood of relapse
- More rapid cycling
- Higher risk of future mixed episodes
- Greater functional impairment
- Higher rates of comorbid conditions such as cardiovascular disease and alcohol/substance use disorder
- Higher risk of suicide

Furthermore, bipolar disorder patients with mixed features may be more likely to experience an **affective switch**, which may also lead to negative health outcomes.^{1,11,12}

The Mixed Features Specifier in Bipolar Disorder

The mixed features specifier (MFS) in the DSM-5 allows for a more specified diagnosis of BP to indicate the presence of concurrent opposing mood symptoms in bipolar patients. The MFS can be applied to current depressive, manic, or hypomanic episodes in BP.¹³




Figure 2. Diagnostic and Statistical Manual of Mental Disorders, 5th ed. American Psychiatric Association; 2013.¹³

Understanding Mixed Episode Symptoms

It is important for nurse practitioners to carefully evaluate patients for mixed presentations, as it may help to provide an accurate diagnosis and ultimately appropriately manage the presenting symptoms. A mixed features episode presents with a combination of mood states. Patients with mixed features may display the following examples of symptoms or symptom combinations¹⁴:

- Depressed mood with irritability
- Socially withdrawn
- Sense of despair and suicide contemplation
- Loss of interest/pleasure and widespread apathy
- Difficulty concentrating and overcrowding thoughts
- Increased/decreased sexual activity
- Difficulty sleeping
- Somatization symptoms such as gastrointestinal disorders and headaches

Understanding and recognizing mixed features in patients with BP is crucial for providing appropriate management over the course of the disease.

References

1. Salvatore G, et al. *J Clin Psychiatry*. 2010;71(11):1488-1501.
2. Möller HJ, et al. *Expert Rev Neurother*. 2004;4(6 Suppl 2):S3-S8.
3. Chakrabarty T, et al. *Curr Psychiatry Rep*. 2020;22(3):15.
4. Phillips ML, et al. *Lancet*. 2013;381(9878):1663-1671.
5. McIntyre RS, et al. *J Affect Disord*. 2015;172:259-264.
6. Grande I, et al. *Lancet*. 2016;387(10027):1561-1572.
7. Vieta E, et al. *Nat Rev Dis Primers*. 2018;4:18008.
8. Young AH, et al. *Comprehensive Men's Mental Health*. Cambridge University Press; 2021:147-158.
9. Vieta E, et al. *J Affect Disord*. 2013;148(1):28-36.
10. Bartoli F, et al. *J Affect Disord*. 2020;276:234-240.
11. Tondo L, et al. *Acta Psychiatr Scand*. 2018;138(3):243-252.
12. Maj M, et al. *Am J Psychiatry*. 2002;159(10):1711-1717.
13. *Diagnostic and Statistical Manual of Mental Disorders*. 5th ed. American Psychiatric Association; 2013.
14. Tavormina G. *Psychiatr Danub*. 2021;33(Suppl 9):6-10.